Subterranean Biology 43: 127–143 (2022)
doi: 10.3897/subtbiol.43.81308
https://subtbiol.pensoft.net

RESEARCH ARTICLE

A peer-reviewed open-access journal
Subterranean Biology
Published by The International Society for Subterranean Biology

Two new troglobitic species of *Iansaoniscus* from Brazilian caves (Crustacea, Isopoda, Pudeoniscidae)

Giovanna Monticelli Cardoso[1], Rafaela Bastos-Pereira[1], Rodrigo Lopes Ferreira[1]

1 *Universidade Federal de Lavras (UFLA), Centro de Estudos em Biologia Subterrânea, Departamento de Ecologia e Conservação, Programa de Pós-graduação em Ecologia Aplicada. Lavras, Minas Gerais, Brazil*

Corresponding author: Giovanna Monticelli Cardoso (gmcardoso.bio@gmail.com)

Academic editor: Matthew L. Niemiller | Received 27 January 2022 | Accepted 19 May 2022 | Published 14 June 2022

http://zoobank.org/AAF4E607-B43C-4797-9903-05BD241F9CF1

Citation: Cardoso GM, Bastos-Pereira R, Ferreira RL (2022) Two new troglobitic species of *Iansaoniscus* from Brazilian caves (Crustacea, Isopoda, Pudeoniscidae). Subterranean Biology 43: 127–143. https://doi.org/10.3897/subtbiol.43.81308

Abstract

Iansaoniscus species are troglobitic terrestrial isopods of the Pudeoniscidae family. In this work two new species are described from caves in the Bahia state, northeastern Brazil; *I. leilae* **sp. nov.** from Toca do Gonçalo cave, in the municipality of Campo Formoso; and *I. paulae* **sp. nov.** from Lapa do Bode cave in the municipality of Itaeté. Additionally, ecological notes and conservation status are provided for both new species.

Keywords

Cave fauna, Neotropics, Oniscidea, terrestrial isopods, troglobites

Introduction

Brazil currently shelters 36 described species of troglobitic oniscidean isopods, and this number has highly enhanced in the last few years (Cardoso et al. 2021, 2022; Campos-Filho et al. 2022). New species have continuously been discovered as new karstic areas are sampled. Styloniscidae stands out among the families with cave-restricted taxa, with 25 species (Fernandes et al. 2019; Cardoso et al. 2021, 2022; Campos-Filho et al. 2022); followed by Scleropactidae with five species (Campos-Filho and Araujo 2011; Campos-Filho et al. 2014, 2020), Philosciidae with four species (Campos-Filho et al. 2019, 2020), and Pudeoniscidae with two species (Campos-Filho et al. 2017).

Copyright *Giovanna Monticelli Cardoso et al.* This is an open access article distributed under the terms of the Creative Commons Attribution License (CC BY 4.0), which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.

Pudeoniscidae present nine species in four genera: *Brasiloniscus* Lemos de Castro, 1973, *Pudeoniscus* Vandel, 1963, *Iansaoniscus* Campos-Filho, Araujo & Taiti, 2017, and *Oxossioniscus* Campos-Filho, Lisboa & Cardoso, 2018 (Vandel 1963; Lemos de Castro 1973; Campos-Filho et al. 2017, 2018). Most species of this family occur in the Brazilian Atlantic Forest, with representatives found associated with leaf litter of cocoa, bromeliads, canopy of *Erythrina fusca* Lour., termite nests, and caves (Campos-Filho et al. 2017, 2018).

To date, *Iansaoniscus* is the only genus of this family with troglobitic species (Campos-Filho et al. 2017). The genus comprises two species: *Iansaoniscus georginae* Campos-Filho, Araujo & Taiti, 2017 from Borboletas cave, municipality of Paripiranga and *Iansaoniscus iraquara* Campos-Filho, Araujo & Taiti, 2017 from Buraco do Cão cave, municipality of Iraquara, both in Bahia state.

The genus is distinguished by the unique shape of cephalon with well-developed antennary lobes and the triangular frontal shield, that is laterally interrupted by antennal grooves; pereonite 1 epimeron with dorsolateral furrow reduced or absent; pleon outline continuous with pereon; telson triangular; uropod protopod subquadrangular; and pleopod exopods with no respiratory structures (Campos-Filho et al. 2017).

In the present work, two new troglobitic species of *Iansaoniscus* are described, both from limestone caves in the Bahia state. Furthermore, ecological notes and conservation status are provided for both new species.

Materials and methods

The new species were manually collected, fixed in 70% ethanol, and taken to the Center of Studies on Subterranean Biology of the Federal University of Lavras (CEBS, UFLA), Lavras, Brazil. Photographs and measurements were taken with ZEISS Axio ZoomV16 stereomicroscope coupled with an Axio Cam 506 Color camera. The specimens were dissected and mounted with Hoyer's medium (Anderson 1954). Illustrations were based on photographs or taken with the aid of a camera lucida on a microscope Leica DM750, and digitally drawn using the software GIMP (v. 2.8) (Montesanto 2015, 2016). Some specimens were dried in alcoholic series for the analysis of dorsal cuticular structures using the scanning electron microscope Hitachi TM4000. The material is deposited in the Collection of Subterranean Invertebrates of Lavras (ISLA), UFLA.

Taxonomy

Family Pudeoniscidae Lemos de Castro, 1973

Genus *Iansaoniscus* Campos-Filho, Araujo & Taiti, 2017

Type species. *Iansaoniscus iraquara* Campos-Filho, Araujo & Taiti, 2017, by original designation.

***Iansaoniscus leilae* sp. nov.**
http://zoobank.org/4F20CBD9-585B-48BF-9C18-019EFF7B5EEF
Figs 1–4

Sclreropactidae sp. nov. [sic!] – Souza-Silva and Ferreira 2016: 9, fig. 3F.

Diagnosis. Cephalon lateral lobes with straight distal margin; frontal shield triangular with frontal margin directed upwards with antennal socket depressed, vertex margin depressed; pereonite 1 epimeron without dorsolateral furrow; pereonite 1 grooved on ventral margin for two thirds of its length, pereonite 2 with well-developed ventral lobe; telson with acute apex; antennula distal article with two lateral and two apical aesthetascs; uropod exopod slightly shorter than endopod, as long as distal margin of telson.

Material examined. ***Holotype*:** Brazil • 1 male; Bahia state, municipality of Campo Formoso, Toca do Gonçalo cave (WGS84 -10.510992, -40.894659); 10th December 2018; RL Ferreira leg.; ISLA 95911. ***Paratypes*:** • 1 male (mounted in slide), 2 females (1 mounted in slide); same data as holotype; ISLA 95912. • 1 male, 1 female; 05th January 2008; same location as holotype; ISLA 95913. • 6 females; 05th January 2008; same location as holotype; ISLA 95914. • 1 male (mounted in slide), 1 female; municipality of Campo Formoso, Toca da Tiquara cave (-10.458611, -40.542222); RL Ferreira leg.; ISLA 95917.

Etymology. The new species is named after Dr. Leila Aparecida Souza, UECE, for her contributions on the systematics of Brazilian subterranean Oniscidea.

Description. Maximum length: male and female, 6 mm. Body outline in lateral view as in Fig. 1A, colorless, eyes absent (Fig. 1A–C). Cephalon lateral lobes with straight distal margin; frontal shield triangular with frontal margin directed upward with antennal socket depressed, vertex margin depressed (Fig. 1B, C). Pereonite 1 epimeron without dorsolateral furrow; pereonite 1 grooved on ventral margin for two thirds of its length, pereonite 2 with well-developed ventral lobe (Fig. 1D–F); pereonites 3–7 with sub-quadrangular epimera. Pleonites 3–5 with epimera sub-quadrangular and directed backwards (Figs 1A, G, 2A). Telson (Figs 1G, 2A) slightly wider than long with concave sides, apex acute. Antennula (Fig. 2B) with three articles, distal article longer than second article bearing two lateral and two apical aesthetascs. Antenna (Fig. 2C) surpassing pereonite 3 when extended backwards, fifth article of peduncle as long as flagellum; flagellum with three articles; second and third articles subequal in length, third article with lateral aesthetascs (Fig. 1H). Mandibles (Fig. 2D, E) molar penicil with 7–8 branches, left mandible with 2+1 penicils, right with 1+1 penicils. Maxillula (Fig. 2F) outer branch with 4 + 5 teeth (two apically cleft); inner branch with two penicils. Maxilla (Fig. 2G) with bilobate apex, outer lobe wider than inner lobe, round, covered with thin setae; inner lobe bearing thick setae. Maxilliped (Fig. 2H) basis rectangular; palp basal article with two setae distinct in length; endite sub-rectangular, long medial seta, distal margin with two teeth. Pereopods 1–7 (Fig. 3A, B) merus and carpus with sparse setae on sternal margin; carpus 1 with longitudinal antennal grooming brush, distal seta apically cleft; dactylus with dactylar organ and



Figure 1. *Iansaoniscus leilae* sp. nov. **A** male holotype, lateral view. Female paratype **B** cephalon, frontal view **C** cephalon, dorsal view **D** pereonites 1 and 2, dorsal view, arrow indicating the *noduli laterales* **E** pereonite 1, ventral view **F** pereonites 2 and 3, ventral view **G** pleonite 5, telson and uropod, dorsal view **H** antennal flagellum **I** uropod. Scale bars: 1 mm (**A**); 500 μm (**B–I**).

ungual setae simple not surpassing outer claw. Uropod (Figs 1G, 2A) protopod longer than wide, with lateral groove; endopod inserted proximally, exopod slightly shorter than endopod, exopod slightly longer than distal margin of telson; proximal margin with lateral groove

Male: Genital papilla (Fig. 3C) with triangular ventral shield and subapical orifices. Pleopod 1 (Fig. 3C) exopod ovoid, margin straight, almost three times wider than long; endopod four times as long as exopod, distal portion slightly bent outwards, apex with small setae. Pleopod 2 exopod (Fig. 3D) triangular bearing one seta on outer margin, endopod flagelliform, almost three times as long as exopod. Pleopod 3–5 exopods as in Fig. 3E–G.

Habitat and conservation issues. The Toca do Gonçalo cave presents ca. 500 meters of mapped conduits. It is inserted at the Una geological group of carbonate rocks from the Caatinga Formation, which comprises relatively young rocks deposited around 600 million years ago (Auler 2019). It is located in the Campo Formoso



Figure 2. *Iansaoniscus leilae* sp. nov. Male holotype **A** telson and uropod, dorsal view **B** antennula **C** antenna **D** left mandible **E** right mandible **F** maxillula **G** maxilla **H** maxilliped. Scale bars: 0.2 mm.

municipality, Bahia state, Northeastern Brazil, within the Caatinga (a semiarid biome) (Fig. 9). The cave presents a single horizontal lenticular-shaped entrance, 2 m high and 8 m wide (Fig. 4A). It has two distinct levels: while the upper level is predominantly dry, the lower level used to be frequently filled with phreatic water (Fig. 4B, C). However, as will be discussed further on, several changes (at both local and regional scales) led to a massive reduction in the water table level, thus, exposing several conduits that used to be inaccessible to men in the past decades (Fig. 4C). The main food sources for both the terrestrial and aquatic fauna seem to be particulate organic matter from the surface, although there are some root mats in the water table level and small guano piles in some areas within the cave.

Culver and Sket (2000) defined hotspots of subterranean biodiversity as caves (or cave systems) with 20 or more cave-restricted species. The Toca do Gonçalo cave shelters 22 troglobitic species and represents one of the three hotspots of subterranean biodiversity known for South America (Souza-Silva and Ferreira 2016; Souza-Silva et al. 2021). Most troglobitic species occurring in this cave remain undescribed, with descriptions already known for only seven of them: the beetle *Coarazuphium caatinga*



Figure 3. *Iansaoniscus leilae* sp. nov. Male holotype **A** pereopod 1 **B** pereopod 7 **C** pleopod 1 **D** pleopod 2 **E** pleopod 3 exopod **F** pleopod 4 exopod **G** pleopod 5 exopod. Scale bars: 0.2 mm.

(Pellegrini & Ferreira, 2014); the cricket *Erebonyx catacumbae* (Mello & Ferreira, 2021); the palpigrade *Allokoenenia canhembora* (Souza & Ferreira, 2022); the centipedes *Cryptops spelaeoraptor* (Ázara & Ferreira, 2014a) and *Newportia spelaea* (Ázara & Ferreira, 2014b); the amphipod *Spelaeogammarus trajanoae* (Koenemann & Holsinger, 2000); and the isopod *Pongycarcinia xyphidiorus* (Messana et al. 2002).

Specimens of *I. leilae* sp. nov. (Fig. 4D, E) were observed at the upper level in the first cave surveys, in the late 1990s. Several specimens were observed, especially near some old bat guano piles. At that moment, most of the lower level was filled with water. Hence, even the upper level was moist due to the influence of the high phreatic level on the whole cave atmosphere. However, local inhabitants use to draw water from the cave (as the region is quite dry) and this practice occurred for decades. This removal occurred manually at the beginning, but a diesel pump (Fig. 4F) was installed inside the cave in the 1980s, drying out previously flooded areas (Prevorčnik et al. 2012; Souza-Silva and Ferreira 2016). It is important to mention that the locals



Figure 4. *Iansaoniscus leilae* sp. nov. habitat **A** Toca do Gonçalo cave entrance **B** cave floor in the inner portion of the cave, where specimens were found **C** water table at the cave lower level **D** adult specimen inside the cave, frontal view **E** adult specimen inside the cave, lateral view **F** diesel pump installed inside the cave in the 70's **G** electric pump installed inside the cave in 2010.

used to remove water from the cave once a week to fill up an external reservoir, hence the water table was persistent. However, in 2010, an electric pump employed by a local farmer for irrigation was installed in the cave (Fig. 4G), leading to a pronounced reduction of at least two meters in the water table inside the cave. Even after the removal of this electric pump (in 2012) the water table reduction continued, since artesian wells were built up outside the cave, so the water demand persisted. During a visit to the cave in August 2013, a major reduction in the water level was observed (approximately 3 m compared to the "pristine" level – observed in the 1990s). In 2013, specimens of *I. leilae* sp. nov. were no longer found in the upper level of the cave (that was extremely dry), being only observed on the moist substrates of the lower

level, even though, in lower densities when compared to the first observations. In December 2018, in a visit to the cave, the whole water table was no longer observed, so previously inaccessible areas were explored, revealing the cave to be much longer than previously thought, and only a few cave-restricted species were observed. From the 22 previously observed troglobitic species, only two (one specimen of *I. leilae* sp. nov., and one springtail) were observed in some moist areas located deep inside the cave in previously flooded areas. The single individual of *I. leilae* sp. nov. was found in a small piece of wood, apparently attracted by the organic compounds. It is important to mention that this species may be seriously threatened, as other cave-restricted species from the Toca do Gonçalo cave. Urgent intervention by the competent environmental agency is needed to ensure the protection of this hotspot of subterranean biodiversity in South America, especially considering that even the most relevant caves can now be destroyed in Brazil (Brasil 2022).

Iansaoniscus paulae sp. nov.

http://zoobank.org/4B0755B7-18B1-404D-B6F0-A6F7A86A4EF4
Figs 5–8

Diagnosis. Cephalon with well-developed lateral lobes with round distal margin; frontal shield rounded, weakly developed; pereonite 1 epimeron without dorsolateral furrow; pereonite 1 and 2 without ventral lobes; antennula distal article with two lateral and one apical aesthetascs; third article longer than second; uropod exopod longer than endopod, longer than distal margin of telson.

Material examined. ***Holotype*:** Brazil • 1 male (mounted in slide); Bahia state, municipality of Itaeté, Lapa do Bode cave (WGS84 -12.938889, -41.062222); 26th July 2007; RL Ferreira leg.; ISLA 95915. ***Paratypes*:** • 5 females; same data as holotype; ISLA 95916.

Etymology. The new species is named after Dr. Paula Beatriz Araujo, UFRGS, for her contribution to the knowledge of Brazilian terrestrial isopods.

Description. Maximum length: male, 3 mm; female 5 mm. Body outline as in Fig. 5A, colorless, vestigial eyes (Fig. 5A, B). Cephalon with well-developed lateral lobes, round distal margin; frontal shield rounded, weakly developed (Fig. 5B). Pereonite 1 epimeron without dorsolateral furrow; pereonites 1 and 2 without ventral lobes (Fig. 5C, D); pereonite 3–7 with subquadrangular epimera. Pleonites 3–5 with epimera subquadrangular and directed backwards (Fig. 5A). Telson (Fig. 6A) slightly wider than long with concave sides, acute apex. Antennula (Fig. 5E) with three articles, distal article longer than second article, with two lateral and one apical aesthetascs. Antenna (Fig. 5F, 6B) surpassing pereonite 3 when extended backwards, fifth article of peduncle as long as flagellum, flagellum with three articles; faint suture between second and third, third article longest, aesthetascs on second and third articles. Mandibles (Fig. 6C, D) with molar penicil of 4 branches, left mandible with 2+1 penicils, right with 1+1 penicils. Maxillula (Fig. 6E) outer branch with 4 + 5 teeth (two apically



Figure 5. *Iansaoniscus paulae* sp. nov. Female paratype **A** lateral view **B** cephalon, dorsal view **C** pereonite 1, dorsal view **D** pereonite 1, ventral view **E** antennula **F** antennal flagellum. Scale bars: 0.5 mm (**A, B, D, E**); 0.3 mm (**C**); 0.1 mm (**E**).

cleft); inner branch with two penicils. Maxilla (Fig. 6F) with bilobate apex, outer lobe wider than inner lobe, rounded and covered with thin setae; inner lobe bearing thick setae. Maxilliped (Fig. 6G) basis rectangular; palp basal article with two setae distinct in length; endite rectangular, long medial seta, distal margin with two teeth. Pereopods 1–7 (Fig. 7A, B) merus and carpus with sparse setae on sternal margin; carpus 1 with longitudinal antennal grooming brush, distal setae cleft; dactylar organ and ungual seta simple not surpassing outer claw. Uropod (Fig. 7C) protopod distal margin and exopod proximal margin with lateral groove; endopod inserted proximally, exopod 1.4× longer than endopod and distinctly surpassing distal margin of telson.

Male: Pleopod 1 (Fig. 7D) exopod ovoid, twice wider than long; endopod almost four times longer than exopod, distal portion slightly bent outwards, apex with small setae. Pleopod 2 exopod (Fig. 7E) triangular bearing one seta on outer margin, endopod flagelliform, almost three times longer than exopod. Pleopod 3–5 exopods as in Fig. 7F–H.

Habitat and conservation issues. The Lapa do Bode cave consists of a dolomite cave with 1,430 meters of horizontal projection, with conduits of labyrinthine pattern. It is inserted in an upper unit of the Una geological group of carbonate rocks from the



Figure 6. *Iansaoniscus paulae* sp. nov. Male holotype **A** telson and uropod, dorsal view **B** antenna **C** left mandible **D** right mandible **E** maxillula **F** maxilla **G** maxilliped. Scale bars: 0.2 mm.

Salitre Formation (Fig. 9) (Auler 2019). This cave presents five entrances (Fig. 8B, C) inserted in a rock wall parallel to the Una River, which flows close to the entrances (Fig. 8A). Although most of the cave is dry, the cave atmosphere is considerably moist due to water ponds in some inner areas. The temperature inside the cave is around 24.5 °C. The cave has been used for touristic purposes for the last decades, and the areas where individuals of *I. paulae* sp. nov. were found (inner portions of the cave, quite distant from the entrances) are also visited by tourists. They were observed walking on moist soil (Fig. 8D) and in some bat guano piles. Although the touristic activities apparently have not been altering intensively the cave features, there is concern that it could impact the species, especially by trampling (by visitors) due to the reduced size of the specimens and absence of structures guiding visitors to walk in specific pathways. The main available food resource is bat guano, that are concentrated in the inner chambers of the cave. It is worth mentioning that the Lapa do Bode cave presents ten cave-restricted species, among springtails, harvestmen, spiders, a beetle, a millipede, a silverfish, a palpigrade and a fish. However, only three of those species are currently described: the carabid beetle *Coarazuphium cessaima* (Gnaspini, Vanin & Godoy, 1998); the pholcid spider *Metagonia diamantina* (Machado, Ferreira & Brescovit, 2011); and the catfish *Rhamdiopsis krugi* (Bockmann & Castro, 2010).



Figure 7. *Iansaoniscus paulae* sp. nov. Male holotype **A** pereopod 1 **B** pereopod 7 **C** uropod **D** pleopod 1 **E** pleopod 2 **F** pleopod 3 exopod **G** pleopod 4 exopod **H** pleopod 5 exopod. Scale bars: 0.2 mm.

Discussion

Taxonomy

The shape of cephalon is a unique feature in *Iansaoniscus* species, however it presents great phenotypic plasticity. *Iansaoniscus leilae* sp. nov. and *I. georginae* show similar patterns such as the frontal shield triangular with frontal margin directed upwards, lateral lobes weakly developed and antennal sockets depressed; while *I. paulae* sp. nov. and *I. iraquara* show well-developed lateral lobes with round margins and frontal shield rounded, weakly developed. The two new species are similar to *I. iraquara* due to pereonites 3 to 7 with subquadrangular epimera, maxillulae outer endite with 4+5 teeth (2 of them apically cleft) and absence of dorsolateral furrow on pereonite 1 epimera. However, while *I. iraquara* and *I. leilae* sp. nov. present ventral lobes on pereonites 1 and 2, such structures are absent in *I. paulae* sp. nov. In *I. georginae* the dorsolateral furrow and lobes are present, but weakly developed, while the maxillulae outer endite presents 4+6 teeth (2 apically cleft). Differently from the other three species, *I. georginae* presents subquadrangular epimera from pereonite 2. Regarding the telson shape, *I. iraquara*, *I. georginae* and *I. paulae* sp. nov. share the condition of a narrowly



Figure 8. *Iansaoniscus paulae* sp. nov. habitat **A** limestone bordering the Paraguaçu river, where the Lapa do Bode cave's entrance is located **B** one of the entrances of Lapa do Bode cave **C** a secondary entrance of Lapa do Bode cave **D** adult specimen inside the cave.

rounded apex, which is acute in *I. leilae* sp. nov. Another difference among the species is that pleopod 1 exopod may present two conditions: ovoid (*I. iraquara*, *I. leilae* sp. nov. and *I. paulae* sp. nov.), and subquadrangular (*I. georginae*). In relation to eyes, such structures are absent in *I. iraquara*, *I. georginae* and *I. leilae* sp. nov., but present and vestigial in *I. paulae* sp. nov. Moreover, the *Iansaoniscus* species present distinct patterns of aesthetascs in the antennula distal article: *I. georginae* (four lateral plus one apical pair), *I. leilae* sp. nov. (two lateral plus two apical), *I. paulae* sp. nov. (two lateral plus one apical) and *I. iraquara* (missing antennule in the description). Additionally,



Figure 9. Geographical distribution of *Iansaoniscus* species, with Brazilian biomes and the karst areas delimited in the state of Bahia.

distinct patterns are observed regarding the number of branches in the mandible molar penicil: *I. iraquara* (5 to 7), *I. georginae* (6 to 7), *I. leilae* sp. nov. (7 to 8), *I. paulae* sp. nov. (4). The proportion of uropods exopod and endopod also varies among the species, with exopod as long as endopod in *I. iraquara*; exopod shorter than endopod in *I. leilae* sp. nov.; and exopod longer than endopod in *I. georginae* and *I. paulae* sp. nov. (being longer in *I. paulae* sp. nov. 1.4×).

Ecology and distribution

Until the discovery of *Iansaoniscus* species in the Caatinga biome (semi-arid region) (Fig. 9), the family was believed to be restricted to the Atlantic Forest (Campos-Filho et al. 2018; Cardoso et al. 2018). The other three genera occur in forests along the coast of Brazil, mainly associated with vegetation and/or leaf litter (Vandel 1963; Lemos de Castro 1973; Campos-Filho et al. 2017). Until then, few collection efforts had been made in more arid areas of Brazil, which conditions are not very suitable for the occurrence of isopods in general. In this sense, *Iansaoniscus*, which are restricted to caves in the only Brazilian semi-arid biome (Caatinga), are supposed to represent a geographical relict.

The xeric conditions of Caatinga currently act as a limiting barrier to the dispersion of many hygrophilic components of fauna, such as isopods (Fernandes et al.

2016). However, Wang et al. (2004) dated speleothems and travertine deposits from the Bahia state and estimated 210,000 years of record of wet periods in northeastern Brazil caused by the displacement of the Intertropical Convergence Zone to the south. The rainforest distribution was affected during these wet periods, which formed a corridor between Amazonian and Atlantic Forests (Por 1992; Behling et al. 2000). Posteriorly, Cruz et al. (2009) observed an anti-phased relationship between precipitation in northeastern Brazil and the rest of tropical South America during the Holocene. The authors analyzed oxygen isotopic records from a speleothem in northeastern Brazil from the last 26,000 years and concluded that a predominantly wet climate dominated the currently semi-arid Caatinga. In this sense, the called south-eastern/north-western route (Bigarella et al. 1975), or the route across the Caatinga formation of eastern Brazil (Ledo and Colli 2017), seem to have favored many biotic exchanges in the past. Along this pathway and under climatic changes, some species or groups of species may have been isolated in moister areas, such as the subterranean conditions of many caves in Caatinga (Polhemus and Ferreira 2018), culminating in current geographical relicts, such as *Allokoenenia* (Palpigradi) (Souza & Ferreira, 2022) and *Iansaoniscus* species.

Acknowledgements

The authors would like to thank the Centro Nacional de Pesquisa e Conservação de Cavernas - CECAV and Instituto Brasileiro de Desenvolvimento e Sustentabilidade - IABS (n°. 006/2021. TCCE ICMBio/Vale (01/2018) for the financial support and scholarship provided to GMC; CNPq (National Council for Scientific and Technological Development) for the productivity scholarship provided to RLF (CNPq n. 308334/2018-3); and the team from the Center of Studies in Subterranean Biology (CEBS/UFLA) for the support in the field trips.

References

Anderson LE (1954) Hoyer's Solution as a Rapid Permanent Mounting Medium for Bryophytes. The Bryologist 57: e242. https://doi.org/10.2307/3240091

Auler AS (2019) Histórico, ocorrência e potencial de cavernas no Brasil. In: Rubbioli E, Menin D, Brandi R (Eds) Cavernas. Atlas do Brasil Subterrâneo. Instituto Chico Mendes de Conservação da Biodiversidade, Brasília 14–51.

Ázara LN, Ferreira RL (2014a) *Cryptops* (Cryptops) *spelaeoraptor* n. sp. a remarkable troglobitic species (Chilopoda: Scolopendromorpha) from Brazil. Zootaxa 826(1): 291–300. https://doi.org/10.11646/zootaxa.3826.1.10

Ázara LN, Ferreira RL (2014b) Two new troglobitic *Newportia* (*Newportia*) from Brazil (Chilopoda: Scolopendromorpha). Zootaxa 3881(3): 267–278. https://doi.org/10.11646/zootaxa.3881.3.5

Behling H, Arz HW, Pätzold J, Wefer G (2000) Late Quaternary vegetational and climate dynamics in northeastern Brazil, inferences from marine core GeoB 3104-1. Quaternary Science Reviews 19: 981–994. https://doi.org/10.1016/S0277-3791(99)00046-3

Bigarella JJ, Andrade-Lima D, Riehs PJ (1975) Considerações a respeito das mudanças paleoambientais na distribuição de algumas espécies vegetais e animais do Brasil. Anais do Simpósio Internacional sobre o Quaternário, 411–464.

Bockmann FA, Castro R (2010) The blind catfish from the caves of Chapada Diamantina, Bahia, Brazil (Siluriformes: Heptapteridae): description, anatomy, phylogenetic relationships, natural history, and biogeography. Neotropical Ichthyology 8: 673–706. https://doi.org/10.1590/S1679-62252010000400001

Brasil (2022) Decreto N° 10.935, de 12 de janeiro de 2022. Dispõe sobre a proteção das cavidades naturais subterrâneas existentes no território nacional. Brasília: Diário Oficial da República Federativa do Brasil. https://www.in.gov.br/web/dou/-/decreto-n-10.935-de-12-de-janeiro-de-2022-373591582 [accessed: 24 january 2022]

Campos-Filho IS, Araujo PB (2011) Two new troglobitic species of Scleropactidae (Crustacea: Isopoda: Oniscidea) from Pará, Brazil. Nauplius 19: 27–39. https://doi.org/10.1590/S0104-64972011000100004

Campos-Filho IS, Araujo PB, Bichuette ME, Trajano E, Taiti S (2014) Terrestrial isopods (Crustacea: Isopoda: Oniscidea) from Brazilian caves. Zoological Journal of the Linnean Society 172(2): 360–425. https://doi.org/10.1111/zoj.12172

Campos-Filho IS, Bichuette ME, Montesanto G, Araujo PB, Taiti S (2017) The first troglobiotic species of the family Pudeoniscidae (Crustacea, Isopoda, Oniscidea), with descriptions of a new genus and two new species. Subterranean Biology 23: 69–84. https://doi.org/10.3897/subtbiol.23.20963

Campos-Filho IS, Lisboa JT, Cardoso GM (2018) A new genus and two new species of Pudeoniscidae Lemos de Castro, 1973 (Crustacea: Isopoda: Oniscidea) from Brazil. Journal of Natural History 52(7–8): 457–482. https://doi.org/10.1080/00222933.2018.1437229

Campos-Filho IS, Fernandes CS, Cardoso GM, Bichuette ME, Aguiar JO, Taiti S (2019) Two new species and new records of terrestrial isopods (Crustacea, Isopoda, Oniscidea) from Brazilian caves. Zootaxa 4564 (2): 422–448. https://doi.org/10.11646/zootaxa.4564.2.6

Campos-Filho IS, Fernandes CS, Cardoso GM, Bichuette ME, Aguiar JO, Taiti S (2020) New species and new records of terrestrial isopods (Crustacea, Isopoda, Oniscidea) of the families Philosciidae and Scleropactidae from Brazilian caves. European Journal of Taxonomy 606: 1–38. https://doi.org/10.5852/ejt.2020.606

Campos-Filho IS, Gallo JS, Gallão JE, Torres DF, Horta L, Carpio-Díaz YM, López-Orozco CM, Borja-Arrieta R, Aguiar JO, Bichuette ME (2022) Unique and fragile diversity emerges from Brazilian caves – two new amphibious species of *Xangoniscus* Campos-Filho, Araujo & Taiti, 2014 (Oniscidea, Styloniscidae) from Serra do Ramalho karst area, state of Bahia, Brazil. Subterranean Biology 42: 1–22. https://doi.org/10.3897/subtbiol.42.75725

Cardoso GM, Campos-Filho IS, Araujo PB (2018) Taxonomic revision of *Brasiloniscus* (Oniscidea, Pudeoniscidae) with description of a new species. European Journal of Taxonomy 434: 1–16. https://doi.org/10.5852/ejt.2018.434

Cardoso GM, Bastos-Pereira R, Souza LA, Ferreira RL (2021) *Chaimowiczia*: a new Iuiuniscinae genus from Brazil (Oniscidea, Synocheta, Styloniscidae) with the description of two new troglobitic species. Subterranean Biology 39: 45–62. https://doi.org/10.3897/subtbiol.39.65305

Cardoso GM, Bastos-Pereira R, Ferreira RL (2022) A new species of *Chaimowiczia* from the karstic Serra do Ramalho plateau, Brazil (Oniscidea, Synocheta, Styloniscidae). Subterranean Biology 42: 139–149. https://doi.org/10.3897/subtbiol.42.80274

Cruz FW, Vuille M, Burn SJ, Wang X, Cheng H, Werners M, Edwards RL, Karmann I, Auler AS, Nguyen H (2009) Orbitally driven east–west antiphasing of South American precipitation. Nature Geoscience 2: 210–214. https://doi.org/10.1038/ngeo444

Culver DC, Sket B (2000) Hotspots of subterranean biodiversity in caves and wells. Journal of Cave and Karst Studies 62: 11–17.

Fernandes CS, Batalha MA, Bichuette ME (2016) Does the cave environment reduce functional diversity? PLoS ONE 11(3): e0151958. https://doi.org/10.1371/journal.pone.0151958

Fernandes CS, Campos-Filho IS, Araujo PB, Bichuette ME (2019) Synopsis of terrestrial isopods (Crustacea: Isopoda: Oniscidea) from Brazilian caves, with emphasis on new records from north, midwest, northeast and southeast regions. Journal of Natural History 53 (17–18): 1095–1129. https://doi.org/10.1080/00222933.2019.1634225

Gnaspini P, Vanin SA, Godoy NM (1998) A new genus of troglobitic carabid beetles from Brazil (Coleoptera, Carabidae, Zuphiini). Papéis Avulsos de Zoologia 40: 297–309.

Koenemann S, Holsinger JR (2000) Revision of the subterranean amphipod genus *Spelaeogammarus* (Bogidiellidae) from Brazil, including descriptions of three new species and considerations of their phylogeny and biogeography. Proceedings of The Biological Society of Washington 113: 104–123.

Ledo RMD, Colli GR (2017) The historical connections between the Amazon and the Atlantic Forest revisited. Journal of Biogeography 44: 2551–2563. https://doi.org/10.1111/jbi.13049

Lemos de Castro A (1973) Pudeoniscidae, família nova, com descrição de um gênero novo e três espécies novas de isópodos terrestres (Isopoda, Oniscoidea). Boletim do Museu Nacional, Nova Serie, Zoologia 287: 1–10.

Machado EO, Ferreira RL, Brescovit AD (2011) A new troglomorphic *Metagonia* Simon, 1893 (Araneae, Pholcidae) from Brazil. Zootaxa 3135(1): 59–62. https://doi.org/10.11646/zootaxa.3135.1.4

Mello FDAG, Ferreira RL (2021) *Erebonyx catacumbae*, n. gen. et sp.: a blind, troglobitic cricket from Brazil (Orthoptera, Grylloidea, Phalangopsidae). Zootaxa 4975(2): 343–356. https://doi.org/10.11646/zootaxa.4975.2.5

Messana G, Baratti M, Benvenuti D (2002) *Pongycarcinia xiphidiourus* n. gen. n. sp., a new Brazilian Calabozoidae (Crustacea Isopoda). Tropical Zoology 15(2): 243–252. http://doi.org/10.1080/03946975.2002.10531178

Montesanto G (2015) A fast GNU method to draw accurate scientific illustrations for taxonomy. Zookeys 515: 191–206. https://doi.org/10.3897/zookeys.515.9459

Montesanto G (2016) Drawing setae: a GNU way for digital scientific illustrations. Nauplius 24: e2016017. https://doi.org/10.1590/2358-2936e2016017

Pellegrini TG, Ferreira RL (2014) Ultrastructural analysis and polymorphisms in *Coarazuphium caatinga* (Coleoptera: Carabidae, Zuphiini), a new Brazilian troglobitic beetle. Zootaxa 3765(6): 526–540. https://doi.org/10.11646/zootaxa.3765.6.2

Polhemus DA, Ferreira RL (2018) Two unusual new genera of cavernicolous Hydrometridae (Insecta: Heteroptera) from eastern Brazil. Tijdschrift voor Entomologie 161(1): 25–38. https://doi.org/10.1163/22119434-00002072

Por FD (1992) Sooretama: The Atlantic Rain Forest of Brazil. The Hague, SPB Academic Publishing, 130 pp.

Prevorčnik S, Ferreira LF, Sket B (2012) Brasileirinidae, a new isopod family (Crustacea: Isopoda) from the cave in Bahia (Brazil) with a discussion on its taxonomic position. Zootaxa 3452: 47–65. https://doi.org/10.11646/zootaxa.3452.1.2

Souza MFVR, Ferreira RL (2022) Two extraordinary troglobitic species of *Allokoenenia* (Eukoeneniidae: Palpigradi) from Brazil: first records of this initially monotypic genus more than a century after its description. European Journal of Taxonomy 789: 11–48. https://doi.org/10.5852/ejt.2022.789.1627

Souza-Silva M, Ferreira RL (2016) The first two hotspots of subterranean biodiversity in South America. Subterranean Biology 19: 1–21. https://doi.org/10.3897/subtbiol.19.8207

Souza-Silva M, Cerqueira RFV, Pellegrini TG, Ferreira RL (2021) Habitat selection of cave-restricted fauna in a new hotspot of subterranean biodiversity in Neotropics. Biodiversity and Conservation 30(14): 4223–4250. http://doi.org/10.1007/s10531-021-02302-8

Vandel A (1963) Isopodes terrestres recueillis en Amérique du Sud par Claude Delamare Debouttéville. In: Debouteville CD, Rapoport EH (Eds) Biologie de l'Amérique Austral, Paris 2: 63–100.

Wang X, Auler AS, Edwards L, Cheng H, Cristalli PS, Smart P, Richards DA, Shen C (2004) Wet periods in northeastern Brazil over the past 210 kyr linked to distant climate anomalies. Letters to Nature 432(9): 740–743. https://doi.org/10.1038/nature03067